

MCQS^{Q&As}

Multiple-choice questions for general practitioner (GP) Doctor

Pass Test Prep MCQS Exam with 100% Guarantee

Free Download Real Questions & Answers **PDF** and **VCE** file from:

<https://www.pass2lead.com/mcqs.html>

100% Passing Guarantee
100% Money Back Assurance

Following Questions and Answers are all new published by Test Prep
Official Exam Center

- ⚙️ **Instant Download** After Purchase
- ⚙️ **100% Money Back** Guarantee
- ⚙️ **365 Days** Free Update
- ⚙️ **800,000+** Satisfied Customers



QUESTION 1

A young boy presented with pain, swelling and redness of right side testes and scrotum. Sudden onset pain relieved on uplifting the scrotum.

What would be the diagnosis?

- A. Torsion testes
- B. Inguinal hernia
- C. Hydrocele
- D. Epididymo-orchitis

Correct Answer: D

According to Prehn's sign, the physical lifting of the testicles relieves the pain of epididymitis but not pain caused by testicular torsion. Negative Prehn's sign indicates no pain relief with lifting the affected testicle, which points towards testicular torsion which is a surgical emergency

QUESTION 2

A 6-year-old girl presents with her 3rd episode of pallor and mildly icteric sclera associated with symptoms of mild upper RTI. On examination, she has an easily tipped spleen.

Which one of following investigations is most likely to confirm her diagnosis?

- A. Eosin-5-maleimide (E5M) staining test
- B. Hb electrophoresis
- C. Howell-jolly bodies in a RBC smear
- D. Reticulocyte count

Correct Answer: B

QUESTION 3

Greatest benefit of thrombolysis is when given in Acute MI?

- A. less than 12 hours from onset of pain
- B. Same benefit in A and B
- C. less than 24 hours from onset of pain

Correct Answer: A

The greatest benefit is less than 12 hours. Very rarely patient gets thrombolytic in 30-60 minutes of chest pain especially

in developing country, no resources to transport, and no good medical care nearby, literature shows advantage of thrombolytic after onset of chest pain is greatest by before 12 hours ...for practical purpose earlier is better may be in 30-60 mins, but you gets mortality reduction advantage if thrombolytic taken within 12 hours of chest pain.

QUESTION 4

Hypothyroid patient on thyroxin had anorexia, dry cough and dyspnea and left ventricular dysfunction. She had normal TSH and T4 levels, Hyperphosphatemia and hypocalcemia.

The diagnosis is:

- A. Hypopituitarism
- B. Primary Hypopituitarism
- C. Secondary Hypopituitarism
- D. Uncontrolled Hypopituitarism

Correct Answer: B

Primary hypoparathyroidism is defined by a low concentration of PTH with a concomitant low calcium level. In pseudo hypoparathyroidism, the serum PTH concentration is elevated as a result of resistance to PTH caused by mutations in the PTH receptor system. In secondary hypoparathyroidism, the serum PTH concentration is low and the serum calcium concentration is elevated.

QUESTION 5

A patient's wife comes to take her husband home after a Myocardial Infarction and asks how long they should wait before they have sex.

What do you tell her?

- A. No waiting necessary
- B. After echocardiography
- C. 2-6 weeks
- D. They should wait for a normal angiography

Correct Answer: C

Some waiting is necessary to have sex after an infarction. Sex minimally increases the risk of infarction. The duration and the intensity of exertion are sufficient to provoke ischemia in some cases