

MCQS^{Q&As}

Multiple-choice questions for general practitioner (GP) Doctor

Pass Test Prep MCQS Exam with 100% Guarantee

Free Download Real Questions & Answers **PDF** and **VCE** file from:

<https://www.pass2lead.com/mcqs.html>

100% Passing Guarantee
100% Money Back Assurance

Following Questions and Answers are all new published by Test Prep
Official Exam Center

- ⚙️ **Instant Download** After Purchase
- ⚙️ **100% Money Back** Guarantee
- ⚙️ **365 Days** Free Update
- ⚙️ **800,000+** Satisfied Customers



QUESTION 1

A 32-year-old woman presents with malaise fatigue and fever. She complains about weight loss. Exam: malar nasolabial fold can be seen.

What is the most appropriate investigation?

- A. Anti-histone
- B. Anti-centromere
- C. Anti-ds DNA
- D. Anti Jo
- E. Anti Scl70

Correct Answer: C

Features are suggestive of SLE. So most appropriate investigation from the given option DNA.

QUESTION 2

The most common cause of SAH (subarachnoid hemorrhage) is _____.

- A. Trauma
- B. Congenital arteriovenous malformation
- C. Berry Aneurysms
- D. HTN

Correct Answer: C

Subarachnoid hemorrhage is sudden bleeding into the subarachnoid space. The most common cause of spontaneous bleeding is a ruptured aneurysm. Symptoms include sudden, severe headache, usually with loss or impairment of consciousness.

QUESTION 3

A 31-year-old woman presents to the ED with a sudden onset of left-sided chest pain and difficulty breathing. She states her only medication is birth control pills. She has smoked 1 pack of cigarettes per day for 10 years. She is tachypneic (24 BrPM) and her heart rate is 120 beats per minute. Physical examination reveals diminished breath sounds on the left and the trachea deviated to the right.

What is the most likely diagnosis?

- A. Tension pneumothorax
- B. Pulmonary embolus

C. Pericardial tamponade

Correct Answer: A

Tension pneumothorax presents with decreased breath sound on one side and tracheal deviation PE does not give tracheal deviation, although it does have chest pain and tachycardia. Muffled heart sounds are seen typically in pericardial tamponade. This patient's risk for pneumothorax is that she is a smoker. It is likely she has a pleural bleb that burst due to her smoking history.

QUESTION 4

Which of the following is the main difference between saphenous vein grafts and internal mammary artery grafts?

- A. There is less need for aspirin and metoprolol with internal mammary artery grafts.
- B. Internal mammary artery grafts remain open for 10 years.
- C. Heparin is necessary for vein grafts.
- D. Warfarin is necessary with saphenous vein grafts.

Correct Answer: B

The main difference between saphenous vein grafts and internal mammary artery grafts is that vein grafts start to become occluded after 5 years but internal mammary artery grafts are often patent at 10 years. There is no difference in the need for medications.

QUESTION 5

A 77-years-old male presents with history of enuresis and change in behavior. Exam: waddling gait.

What is the most likely diagnosis?

- A. Subdural haemorrhage
- B. Normal pressure hydrocephalus
- C. Psychotic depression
- D. Brain tumor

Correct Answer: B

[Latest MCQS Dumps](#)

[MCQS VCE Dumps](#)

[MCQS Study Guide](#)