

# NCLEX-PN<sup>Q&As</sup>

National Council Licensure Examination(NCLEX-PN)

**Pass NCLEX NCLEX-PN Exam with 100% Guarantee**

Free Download Real Questions & Answers **PDF** and **VCE** file from:

<https://www.pass2lead.com/nclex-pn.html>

100% Passing Guarantee  
100% Money Back Assurance

Following Questions and Answers are all new published by NCLEX  
Official Exam Center

- ⚙️ **Instant Download** After Purchase
- ⚙️ **100% Money Back** Guarantee
- ⚙️ **365 Days** Free Update
- ⚙️ **800,000+** Satisfied Customers



**QUESTION 1**

The vast majority of deaths resulting from unintentional poisoning occur in \_\_\_\_\_.

- A. infants
- B. toddlers
- C. teens
- D. adults

Correct Answer: B

The vast majority of deaths resulting from unintentional poisoning occur in toddlers.

---

**QUESTION 2**

Multifetal pregnancies with triplets occur at a rate of 1 in 8,100 births, but twins occur much more frequently with a rate of \_\_\_\_\_.

- A. 1 in 85 births
- B. 1 in 5400 births
- C. 1 in 2700 births
- D. 1 in 540 births

Correct Answer: A

Twins, whether monozygotic or dizygotic, occur at the rate of about 1 in every 85 births.

This rate has seen increases due to various reasons, such as the frequency of fertility treatments and women waiting until later in life to have children; older women, defined as over 35 years old, are more likely to carry multiples than younger women.

---

**QUESTION 3**

The factor that most determines drug distribution is \_\_\_\_\_.

- A. vascular perfusion of the tissue or organ
- B. salt form
- C. drug interactions
- D. steady state

Correct Answer: A

Drugs are distributed via the circulatory system.

Adequate perfusion is necessary for distribution of a drug.

Salt form, drug interactions and steady state are not as dependent on adequate perfusion.

---

#### QUESTION 4

Paula is a 32-year-old woman seeking evaluation and treatment of major depressive symptoms.

A major nursing priority during the assessment process includes which of the following?

- A. meaning of current stressors
- B. possibility of self-harm
- C. motivation to participate in treatment
- D. presence of alcohol or other drug use

Correct Answer: B

Unless the client is first assessed for self-harm or suicide potential, the staff might not observe the necessary degree of vigilance needed in the client's environment.

Physical needs are the second most critical concern with a depressive client.

Though the client may be encouraged to attend group therapy as part of the treatment plan, the client's safety takes precedence.

Response to medication takes time and is not an initial concern.

---

#### QUESTION 5

Your pediatric weighs 15.8 kg.

How many pounds does this child weigh?

- A. 36 pounds
- B. 33.6 pounds
- C. 35 pounds
- D. 34.8 pounds

Correct Answer: D

Since there are 2.2 pound in one kilogram, you calculate the weight in pounds by multiplying 2.2 by the kg weight as

---

below.

[Latest NCLEX-PN Dumps](#)

[NCLEX-PN Practice Test](#)

[NCLEX-PN Exam Questions](#)